

INDICATIONS FOR ADMISSION TO A HOSPITAL WARD

- Children who have sustained a head injury should be admitted to hospital if any of the following risk factors apply:
 - any indication for a CT scan
 - suspicion of non-accidental injury
 - significant medical comorbidity
 - difficulty making a full assessment
 - child not accompanied by a responsible adult
 - social circumstances considered unsuitable.

- In injured children, especially the very young, the possibility of non-accidental injury must be considered:
 - when findings are not consistent with the explanation given
 - if the history changes, or
 - if the child is known to be on the Child Protection Register.In such cases a specialist paediatrician with responsibility for child protection should be involved. Child protection procedures should be followed.

- Primary and secondary care information systems should identify children on the Child Protection Register and frequent attenders.

INDICATIONS FOR DISCHARGE

- Children can be discharged from the ED if no additional risk factors apply.

REFERRAL TO NEUROSURGICAL UNIT

- D** Features suggesting that specialist neuroscience assessment, monitoring, or management are appropriate include:
- persisting coma (GCS score 8/15 or less) after initial resuscitation**
 - confusion which persists for more than four hours**
 - deterioration in level of consciousness after admission** (a sustained drop of one point on the motor or verbal subscales, or two points on the eye opening subscale of the GCS)
 - focal neurological signs**
 - a seizure without full recovery**
 - compound depressed skull fracture**
 - definite or suspected penetrating injury**
 - a CSF leak or other sign of a basal fracture.**

TRANSFER TO A NEUROSURGICAL UNIT

- Transfer of a child to a specialist neurosurgical unit should be undertaken by staff experienced in the transfer of ill children, such as the Scottish Paediatric Retrieval Service.
- Consultation on the best method of transfer of an individual patient should be with referring healthcare professionals, transfer clinicians and receiving neurosurgeon. It should take into account the clinical circumstances, skill of available staff, imaging, mode of transfer and timing issues.

DISCHARGE PLANNING AND ADVICE

- Before discharge from the ward a patient with a head injury must be assessed by an experienced doctor, who must establish that all the following criteria have been met:
 - consciousness has recovered fully and is sustained at the pre-injury state
 - the patient is eating and drinking normally and not vomiting
 - neurological symptoms/signs have either resolved, or are minor and resolving or are amenable to simple advice/treatment, (eg headache relieved by simple analgesia, or momentary positional vertigo due to vestibular disturbance)
 - the patient is either mobile and self caring or returning to a safe environment with suitable social support
 - the results of imaging and other investigations have been reviewed and no further investigation is required
 - extracranial injury has been excluded or treated.
- Clear written instruction should be given to and discussed with parents or carers before a child is discharged.

FOLLOW UP

- Children suffering from moderate/severe head injury should be followed up by a specialist multidisciplinary team to assess rehabilitation needs.
- Parents should be given information and advice about the possible short/longer term difficulties that their child may have.
- The primary healthcare team, school health team and teachers should be notified of all children with head injury regardless of severity.

This Quick Reference Guide provides a summary of the main recommendations relating to children in **SIGN guideline 110: Early management of patients with a head injury.**

Recommendations are graded **A B C D** to indicate the strength of the supporting evidence. Good practice points are provided where the guideline development group wishes to highlight specific aspects of accepted clinical practice.

Details of the evidence supporting these recommendations can be found in the full guideline, available on the SIGN website: www.sign.ac.uk



Early management of children with a head injury
Quick Reference Guide

May 2009

ASSESSMENT AND CLASSIFICATION

D The management of patients with a head injury should be guided by clinical assessments and protocols based on the Glasgow Coma Scale and Score.

Great care should be taken when interpreting the Glasgow Coma Scale in the under fives and this should be done by those with experience in the management of the young child.

THE PAEDIATRIC COMA SCALE AND SCORE

FEATURE	SCALE RESPONSES	SCORE
Eye opening	Spontaneous	4
	To voice	3
	To pain	2
	None	1
Verbal response	Orientated/interacts/follows objects/ smiles/alert/coos/babbles words to usual ability	5
	Confused /consolable	4
	Inappropriate words/ moaning	3
	Incomprehensible sounds/irritable/ inconsolable	2
	None	1
Best motor response	Obeys commands/ normal movement	6
	Localise pain/ withdraws to touch	5
	Withdrawal to pain	4
	Flexion to pain	3
	Extension to pain	2
	None	1

TOTAL COMA 'SCORE' 3/15 – 15/15

INDICATIONS FOR REFERRAL TO THE ED

B Children with any of the following signs and symptoms should be referred to an appropriate hospital for further investigation of potential brain injury:

- GCS < 15 at initial assessment
- post-traumatic seizure (*generalised or focal*)
- focal neurological signs
- signs of a skull fracture
- loss of consciousness
- severe and persistent headache
- repeated vomiting (*two or more occasions*)
- post-traumatic amnesia > 5 minutes
- retrograde amnesia > 30 minutes
- high risk mechanism of injury
- coagulopathy, whether drug-induced or otherwise
- clinical suspicion of non-accidental injury.

-
- significant medical comorbidity
 - difficulty making a full assessment
 - not accompanied by a responsible adult
 - social circumstances considered unsuitable.

INDICATIONS FOR HEAD CT

